

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> A31N 1/02, 63/00, A61B 19/00	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/17514</b> <b>(43) International Publication Date:</b> 13 June 1996 (13.06.96)
<b>(21) International Application Number:</b> PCT/US95/15738 <b>(22) International Filing Date:</b> 5 December 1995 (05.12.95)  <b>(30) Priority Data:</b> 08/349,747      5 December 1994 (05.12.94)      US  <b>(71) Applicant:</b> NEW YORK BLOOD CENTER, INC. [US/US]; 310 East 67th Street, New York, NY 10021 (US).  <b>(72) Inventors:</b> RUBINSTEIN, Pablo; 92 Woodhollow Lane, New Rochelle, NY 10804 (US). COELHO, Philip, Henry; 121 Giotto Way, El Dorado Hills, CA 95762 (US). STEVENS, Cladd, E.; Apartment W8F, 420 East 64th Street, New York, NY 10021 (US).  <b>(74) Agents:</b> GEORGE, Kenneth, P. et al.; Amster, Rothstein & Ebenstein, 90 Park Avenue, New York, NY 10016 (US).		<b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> METHOD AND BAG SET FOR CONCENTRATING WHITE CELLS  <b>(57) Abstract</b>  A therapeutic product formed from a high concentration of white blood cells having a high degree of cell viability. The white blood cells are sequestered from their normal population presence in whole blood by placing the blood into a container and preventing coagulation of the blood, separating the blood into two components, one of which is extremely rich in white blood cells through the use of a reagent and centrifugation, sequestering the white cell concentration, and freezing the white cells.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/15738

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claims 1-24, drawn to an apparatus for forming the product, the product and a method of using the apparatus to make the product.

Group II, claims 25-39, drawn to drawn to a second process.

The inventions listed as Groups I and II do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The process of group II is not related so as to form a single inventive concept because the bag system (apparatus) of Group I used in the process of Group I to produce the product of group I is not required in the process of Group II. Multiple processes are not permissible in a single application under Rule 13.1.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/15738

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A01N 1/02, 63/00; A61B 19/00  
US CL : 435/2; 424/93.71; 604/410, 416

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/2; 424/93.71; 604/410, 416

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MEDLINE, BIOSIS, APS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	US, A, 4,004,975 (LIONETTI et al.) 25 January 1977, see entire document, particularly Figure 1.	1-20 ----- 21-39
X -- Y	US, A, 5,004,681 (BOYSE et al.) 02 April 1991, see column 41, lines 25-35, column 40, lines 9-30.	18-24 ----- 25-39
X -- Y	US, A, 4,332,122 (WILLIAMS) 01 January 1982, see entire article.	1 ----- 2-39

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

•	Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance		
"E"	earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed	"&"	document member of the same patent family

Date of the actual completion of the international search

20 MARCH 1996

Date of mailing of the international search report

27 MAR 1996

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SANDRA SAUCIER

Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US95/15738

## Box I (Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet))

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☒ No protest accompanied the payment of additional search fees.

METHOD AND BAG SET FOR CONCENTRATING WHITE CELLS

5

FIELD OF THE INVENTION

The following invention is directed generally to the therapeutic utilization of white blood cells, a technique for sequestering the white blood cells by causing them to coalesce in a population density greater than they normally occur in nature, and a method for causing an enriched concentration in conjunction with an array of bags oriented in a set that facilitates both the concentration process and a method for preserving the white blood cells.

BACKGROUND OF THE INVENTION

It is now recognized that placenta/umbilical cord blood (PB) contains large numbers of hematopoietic stem and progenitor cells that endow PB with extraordinary therapeutic capabilities in the reconstitution of bone marrow damaged as a result of inherited diseases, accidents or medical procedures. As in the case of ordinary collection of bone marrow for transplantation, PB contains immune cells potentially capable of mounting specific responses against the recipients of such transplants, but in contrast to adult immunological cells, those in PB display a lower, perhaps much lower tendency to produce damaging immune responses against the recipient. The clinical syndrome produced by the immune responses of the graft against the recipient's cells and tissues is designated "Graft versus Host Disease" (GVHD). In the typical clinical situation, the recipient's own immune response against the graft is abrogated by drugs and irradiation treatments designed to reduce or eliminate the immunological and other hematopoietic cells and thus avoid the host versus graft immune reaction that would cause rejection of the graft. It has been proven that the principal targets of these Graft versus Host and Host versus Graft immune reactions are antigens encoded by the

-2-

genes of the HLA (Human Leukocyte Antigen) system and that successful outcomes of bone marrow transplants are dependent on the sharing of HLA antigens by donor and recipient. Sibling donors who have inherited the same paternal and maternal HLA genes present in the recipient are HLA-identical and thus, optimal from this viewpoint. Patients lacking such HLA-identical sibling donors must receive transplants from more distant relatives or from unrelated donors. Because the HLA system includes several discrete genes each of which displays an extremely large number of antigenically different variants in the population, such distant relative-donor or unrelated-donor transplants must be expected to contain a variable number of HLA incompatibilities unless they are selected from among potential donors by identifying the specific variants present in each and choosing donors whose HLA antigens match those of the recipient. To perform this selection with significant probability of success, it is necessary to have access to large panels of potential donors whose HLA antigens are known. In the case of unrelated donor PB, this requires establishing a bank of frozen HLA-typed units collected from random placentas. Heretofore, the most widely accepted method for freezing PB consisted of adding to the whole PB unit an equal volume of a cryopreservative solution, with the double disadvantage that the volume of each cryopreserved unit becomes very large and that a relatively large amount of possibly deleterious cryopreservative is eventually administered to the recipients of such PB units. Administration of cryoprotectant and hemoglobin from erythrocytes destroyed by using a freezing and thawing method designed to protect the stem and progenitor cells but not the erythrocytes may have toxic effects generally and especially on specific organs such as the kidney of the recipient. In addition, there is the logistical consequence that a large number of freezers would be needed to contain useful numbers of the large volume



-3-

frozen units in reserve, with the attending increase in up-front and running costs. The applicants have developed a practical method that allows a substantial reduction of the volume of PB Units by eliminating the unneeded mature red blood cells and an equivalent volume of plasma. This submission describes this method and a specially designed set of plastic bags and connecting tubes intended to facilitate the accomplishment of the desired concentration of the needed stem cells and progenitor cells with minimal manipulation and risk of contamination. Essentially, this method will allow an experimental, time consuming laboratory process to become a routine procedure in blood banks.

The following submission reflects the state of the art of which applicant is aware insofar as these documents appear germane to the patent process. However, it is respectfully stipulated that none of these patents teach singly nor render obvious when considered in any conceivable combination the nexus of the instant invention as set forth hereinafter.

	<u>INVENTOR</u>	<u>PATENT NO.</u>	<u>ISSUE DATE</u>
	Tenczar, Jr.	3,187,750	06/1965
25	Williams	4,332,122	06/1982
	Pattillo, et al.	4,937,194	06/1990
	Boyse, et al.	5,004,681	04/1991
	Carmen, et al.	5,104,788	04/1992
	Bauman, et al.	5,154,716	10/1992
30	Boyse, et al.	5,192,553	03/1993

OTHER PRIOR ART (Including Author, Title, Date, Pertinent Pages Etc.) Pablo Rubinstein, Richard E. Rosenfield, John W. Adamson and Cladd E. Stevens (The Lindsley F. Kimball Research Institute of The New York Blood Center); Stored Placental Blood for Unrelated Bone Marrow Reconstitution; May, 1993; entire paper.

-4-

SUMMARY OF THE INVENTION

The ~~therapeutic product~~ of the present invention is advantageous, first, because it recovers all or almost all of the stem and progenitor cells of the original collection of PB in a small and uniform volume that  
5 requires minimal and predictable storage space, second, because it permits a consistent methodology for processing PB units which results in a routinely dependable product with less dependence on operator skill and third, because  
10 the potentially deleterious effects of the cryoprotectant and of the free hemoglobin are minimized.

One first aspect to the nature of the product improved according to the present invention involves the methodology by which the white blood cells (which include  
15 the hematopoietic stem and progenitor cells) are separated from the bulk of other components in the whole PB and the manner in which the viability of such white cells is preserved by avoiding exposure to bacterial and fungal contamination, potentially damaging chemical agents,  
20 excessive centrifugal forces and osmotic imbalances. Typically, bacterial and/or fungal contamination occurs when PB or white blood cell suspensions derived from PB are exposed to ambient air in the course of preparatory manipulations; chemical damage is possible when certain  
25 chemicals are used to lyse the accompanying red blood cells or to aggregate white cells; and physical damage follows the use of excessive centrifugal speed in separation of the cellular components of the blood according to their density, by centrifugal stratification.  
30 In addition, the method according to the present invention provides for avoidance of prolonged exposure of the separated white blood cells to cryopreservation solutions at room temperature, an exposure that results in decreased viability of the white blood cells and of the stem and  
35 progenitor cells contained therein because of osmotic imbalances and, possibly, other toxic effects of the intracellular cryoprotectants themselves.

-5-

Another aspect of the present invention involves the set of interconnected plastic containers (designated as bags). The set under the present invention permits a selective concentration of the white blood cells and of the stem and progenitor cells contained therein without reducing their normally high viability and freedom from contamination by infectious organisms from the environment. Whole PB is collected into a mother bag and is subsequently processed through a series of bags of appropriate chemical structure and physical shape and capacity culminating in storage of a separated fraction containing most of the white blood cells of the collected PB in liquid nitrogen at -196 C inside a specially constructed freezer bag. Intervening steps include the addition of substances that enhance the aggregability of red blood cells and the separation of components by transferring supernatants into connected satellite bags. A special bag and its connecting assembly permits the addition of measured amounts of cryoprotectant to the separated white blood cell concentrate. This connecting assembly allows the cryoprotectant to be added to the white cells at a precise, slow speed required to maintain optimal cell viability.

The bag which is to be used for freezing and storage includes a plurality of connected, but detachable compartments for sequestration of the white blood cells into different discrete chambers. One chamber, the main compartment, is intended to keep the bulk of the white blood cells. A smaller compartment lends itself to the storage of a smaller fraction of the bag contents which may be separated from the main compartment without thawing, and extemporaneously detached from it for separate thawing and subsequent in vitro expansion of the hematopoietic stem and progenitor cell populations contained in the corresponding fraction of the white blood cells. A third and subsequent chamber contains very small aliquots of the white blood cell suspension and are

-6-

intended to serve as detachable samples for testing the aptness of the unit to be transplanted or assessing its suitability as donor tissue for a specific recipient. The freezing bag also includes indicia on the outer surface of each of its detachable areas for identification of the specific unit that will be stored in it, to facilitate storage and retrieval from designated sectors of cryogenic storage depots. Means are also provided in an exterior surface of the freezer bag to facilitate the placing and removing of the freezer bag into and from, respectively, its assigned storage location by automated instrumentation.

#### OBJECTS OF THE INVENTION

Accordingly, it is a primary object of the present invention to provide the means for preparing PB derived hematopoietic stem and progenitor cells in a novel and therapeutically more useful form. The product becomes a bag containing a high concentration of white blood cells having a high degree of cell viability.

A further object of the present invention is to provide a novel and useful method for generating the therapeutic product according to the previous object.

A further object of the present invention is to provide an aseptic and interconnected bag set for use in conjunction with the method of developing the therapeutic product herein above.

A further object of the present invention is to provide a freezer storage bag configured to contain the therapeutic dose in a cryoprotected environment for protracted periods of time until needed for dosage.

A further object of the present invention is to provide a freezer bag as noted above provided with a plurality of compartments in which the therapeutic dose has been sequestered so that various aliquots can be strategically excised from the freezer bag for several purposes.

-7-

Viewed from a first vantage point, it is an object of the present invention to provide a system for developing placental stem cells, comprising in combination: a first blood bag adapted to receive blood  
5 from a placenta therewithin, means within the blood bag to prevent coagulation, reagent means removably coupled to the blood bag, means for separating supernatant from the first blood bag and into a white cell bag, means for separating white cells from plasma in the white cell bag,  
10 a plasma bag removably coupled to the white cell bag for receiving the expressed plasma from the white cell bag, cryoprotectant means operatively coupled to the white cell bag, and a stem cell freezing bag operatively coupled to the white cell bag for transferring contents from the  
15 white cell bag to the stem cell freezing bag.

Viewed from a second vantage point, it is an object of the present invention to provide a method for preparing concentrated and partially purified white blood cell suspensions containing placental stem cells,  
20 comprising the steps of: placing blood from a placenta into a first blood bag, preventing coagulation within the blood bag, coupling reagent means into the blood bag, centrifuging and separating white blood cell rich supernatant from the first blood bag and placing the  
25 supernatant into a white cell bag, separating white cells from plasma in the white cell bag, removably coupling a plasma bag to the white cell bag and expressing the plasma from the white cell bag into the plasma bag. Coupling cryoprotectant means to the white cell bag, transferring  
30 contents from the white cell bag to a stem cell freezing bag, and freezing the stem cell freezing bag with its contents follows.

Viewed from a third vantage point, it is an object of the present invention to provide a therapeutic  
35 product comprising at least 80% of the white blood cells (including stem and progenitor cells) with viability greater than 90% and fewer than 10% of the red blood cells

-8-

in the original PB collection.

These and other objects will be made manifest when considering the following detailed specification when taken in conjunction with the appended drawing figures.

5

#### BRIEF DESCRIPTION OF THE DRAWING FIGURES

Figure 1 is a schematic view of the stem cell processing bag set according to the present invention.

10 Figure 2 is a detailed view of the freezing bag shown in figure 1.

Figure 3 is a view similar to figure 2 showing the interior of the freezing bag.

Figure 4 is a flow chart the method according to the present invention.

15

#### DESCRIPTION OF PREFERRED EMBODIMENTS

Referring to the drawings now, wherein like reference numerals refer to like parts throughout various figures, reference numeral 100 is directed to an apparatus  
20 according to the present invention.

In essence, the apparatus 100 may be viewed as three arrays of bags collectively defining a bag set. Individual bags are provided with removable connection means to assure selected admission into the several bags  
25 only under aseptic conditions. In a preferred form of the invention, the array of bags 100 includes six bags: a blood bag 10 defining a first array; a reagent bag 20, a white cell bag 30, a plasma bag 40 and a cryoprotectant bag 50 defining a second array; and a stem cell freezing  
30 bag 60 defining a third array. Cord blood (i.e. blood from the placenta and umbilical cord) is admitted to the blood bag 10 which had previously been dosed with an anticoagulant. Next, the second array is connected to the blood bag 10. A separation reagent is admitted to the  
35 blood bag via conduit 26 from reagent bag 20. Centrifuging blood bag 10 follows. Supernatant containing the white blood cells is expressed off into the white cell

-9-

bag 30, whereupon further centrifugation takes place. Next, supernatant plasma is expressed off into the plasma bag 40 leaving sedimented white blood cells in white cell bag 30. Cryoprotectant from the cryoprotectant bag 50 is transferred via conduit 59 to the white cell bag 30 slowly. Subsequently, the contents of the white cell bag 30 are transferred to the stem cell freezing bag 60 which is thereafter frozen and stored in liquid nitrogen for subsequent use.

More specifically, and with reference to figure 1, whole, placental, and umbilical cord blood is collected into a blood bag 10 provided with an anticoagulant such as Citrate, Phosphate and Dextrose (CPD). Assume, for the sake of explanation, that one hundred (100) milliliters of blood are placed within the blood bag. Typically, cord blood will exhibit a ratio of one thousand (1,000) red cells to each "non-red" cell (for simplicity, assume the non-red blood cell can be labeled white blood cells). Naturally, the main recognizable and functionally capable cells circulating in blood include erythrocytes, neutrophilic, eosinophilic, and basophilic granulocytes; B- T- and non B- non T-lymphocytes; monocytes and platelets. These mature cells derive from and are replaced, on demand, by morphologically recognizable dividing precursor cells for the respective lineages such as erythroblasts for the erythrocyte series, myeloblasts, promyelocytes and myelocytes for the granulocyte series, and megakaryocytes for the platelets. The precursor cells derive from more primitive cells that can simplistically be divided into two major subgroups: stem cells and progenitor cells. Of course, neonatal blood has other cellular constituents which will not be discussed here so as not to obscure the essence of the invention. The blood bag 10 includes at least two access portals. A first portal 2 receives the cord blood whereupon the access portal 2 is sealed. Typically sealing includes a heat seal to insure asepsis. A second portal 4 is provided

-10-

which communicates with a spike 22 coupled via conduit 26 to a separation reagent bag 20 and through conduit 32 to the white cell bag 30 from the second array of bags discussed above. In addition, the blood bag 10 may also  
5 be provided with a third access 6 which may include a sample tube, should it be found desirable to place into storage an exemplar of the cord blood which was originally drawn. Access 6 may also provide alternative connections to bag 10.

10           Once the cord blood has been admitted into the blood bag 10, the admixture with an anticoagulant such as CPD prevents the clotting of the placental blood and readies the blood for admixture with a reagent contained within reagent bag 20. After the admission of the reagent  
15 to the blood bag and thorough mixing, the bag is centrifuged at a precise speed and the white-cell-rich supernatant is expressed into the white cell bag 30. The reagent is intended to facilitate the sedimentation of the red blood cells which is greatly accelerated by a very  
20 light centrifugation step ( $50 \times G \times 5 \text{ min.}$ ). The effects of the addition of separation reagent and centrifugation are to produce a supernatant which contains eighty to ninety-five percent (80-95%) of the white blood cells and less than ten percent (10%) of the red blood cells of the  
25 originally collected blood. This reduces the presence of red cells (compared to white cells) by approximately ninety percent (90%). In the white cell bag, the red cell to white cell count ratio is now reduced to approximately one hundred (100) to one (1).

30           Typically, reagents which promote effective separation of the red blood cells from the white blood cells operate on the basis of mechanisms which can be the subject matter of some speculation as to the physical process or model that describes the separation process.  
35 One vantage point advances the premise that the addition of the reagent raises the dielectric strength of the suspension medium and then, its charge-dissipating



-11-

capacity, so that the tendency for the red blood cells to remain in uniform suspension is disturbed. Another view is that the polymeric molecule of the reagent binds to two or more red blood cells, causing them to aggregate and form characteristic "rouleau" i.e., loose clumps of red blood cells stacked together by the flat aspects of their discoidal surface. The effect, however, irrespective of the physical model that one envisions, is that separation between the red and white cells is possible with relatively minor, gentle and brief centrifugation. This accelerates the settling of the red cells and preserves the white cells in the suspended, unmodified state. In a preferred embodiment, once the reagent from bag 20 has been placed within the blood bag 10, centrifugation at fifty (50) gs for approximately five (5) minutes provides effective separation.

Reagents which change the charge dissipation characteristic or alter the dielectric strength of the constituent components can be selected from a relatively broad range of suitable substances. A six percent (6%) concentration of Heptastarch is presently preferred both due to efficacy, cost, and wide spread utilization in clinical blood processing. However, similar natural polymers such as dextrans, gelatins, modified or unmodified starches or synthetics such as polyethylene glycol or polyvinyl-pirrolidone and many others could conceivably be substituted as conditions warrant. A similar effect may also be obtained with substances whose molecules attach with high avidity to two or more red cells such as antibodies and lectins. In any event, any one of these red cell-cryoprecipitating reagents contained in the reagent bag is dispensed from the reagent bag 20 via outlet 24 through branch passageway 26 and through the outlet spike 22 received by portal 4 or, alternatively, portal 6, into the blood bag 10.

Mixing of the reagent with the blood in the bag 10 followed by gentle centrifugation results in a

-12-

separation in which the supernatant composed of plasma, most of the white blood cells and a small fraction of the red blood cells, is expressed off into the white cell bag 30 via a branch passageway 32 communicating between the spike 22 and the bag 30 with a T-adapter 34 which allows a bifurcation between the branch 26 and the branch 32. The bulk of the red blood cells remain in bag 10.

The enriched white cell mixture is prevented from entering the reagent bag by means of a clamp 28 operatively engaged on the branch passage 26. The enriched white cell mixture in the white cell bag 30 at inlet 36 is now ready for further processing.

As an example, assume one hundred (100) milliliters of PB had been originally collected into blood bag 10. A preferred embodiment provides a reagent bag 20 with a sufficient volume of Hydroxyethyl starch (Heptastarch, Dupont) to provide for the addition of a volume equal to one-fifth ( $1/5$ ) that of the PB collection into bag 10. In this example, one-fifth ( $1/5$ ) of one hundred (100) milliliters equals twenty (20) milliliters. Typically, seventy (70) milliliters of white cell enriched supernatant plasma (containing the reagent solution) will be produced which will be expressed into the white cell bag 30. Once there, the contents are subjected to further centrifugation at-four hundred (400) x G x ten (10) minutes. Typically, of the seventy (70) milliliters that had been admitted into the white cell bag 30, fifty-five (55) milliliters will be expressed off thereafter into a plasma bag 40, leaving approximately fifteen (15) milliliters of highly-enriched white cell product in bag 30. The supernatant transferred to bag 40 contains the bulk of the plasma, anticoagulant and reagent and essentially no cells.

The white cell bag 30 includes an outlet portal 38 that communicates with the plasma bag 40 via a branch conduit 42 having a T-adapter 44 and a constrictor 46 in line. The supernatant is expressed from white cell bag 30

-13-

via conduit 42 to the plasma bag 40 via its own portal 48. Once the supernatant has been received into the plasma bag 40 it is sealed off and the plasma bag 40 is disconnected from the white cell bag 30.

5 Cryoprotectant from cryoprotectant bag 50 is next admitted into the white cell bag 30. Cryoprotectant bag 50 includes an outlet 52, a branch passageway 54 and a constrictor element 56 on the line 54 in fluidic communication with the portal 38 of the white cell bag 30  
10 through T-adaptor 44. Typically, three point eight (3.8) milliliters of cryoprotectant is admitted into the fifteen (15) milliliters contained within the white cell bag 30. It is extremely desirable to admit the cryoprotectant into the white cell bag 30 at a relatively slow rate.  
15 Typically, the three point eight (3.8) milliliters of cryoprotectant is admitted into the bag over a twenty (20), minute interval, while continuously mixing the cryoprotectant with the contents of the white cell bag by hand or with an orbital shaker. A preferred  
20 cryoprotectant solution includes Dimethyl Sulfoxide DMSO (an intracellular cryoprotectant) diluted to fifty percent (50%) with dextran an extracellular cryoprotectant. One feature of the instant invention is that the constrictor element 56 determines that the intracellular  
25 cryoprotectant can only enter white cell bag 30 very slowly. Thus, the intracellular cryoprotectant increases its concentration and permeates the white cell mixture contained within the white cell bag 30 without causing damage to the cells. In order to effect same, a metering  
30 instrumentality 58 may be interposed in the branch 54 instead of the constrictor element 56 (should the constriction not provide a constant flow rate) and in fluid communication with the portal 38. The metering instrumentality 58 can be a pump. Alternatively the  
35 cryoprotectant bag and pump arrangement can be replaced with a syringe or other metering apparatus which facilitates the slow addition of cryoprotectant to the

-14-

white cell bag 30.

The physical analogy for the cryoprotectant is that the DMSO penetrates through the white cell membrane and reduces the capacity of intracellular water as it  
5 freezes to crystallize intracellularly and inflict damage to the cell walls. Dextran and other extracellular cryoprotectants such as diverse kinds of soluble starches, proteins and sugars are believed to provide extracellular layers around white cells that insulate the cells from the  
10 tendency of the water to form crystals during the freezing process and to develop excessive extracellular hyperosmolarity, both of which might reduce cell wall integrity and cellular viability. By providing the cryoprotectant at a measured rate, over a relatively long  
15 period of time, cell viability will have been maximized by providing ample time for the DMSO to diffuse into cells and to reach equilibrium across the cell membrane and for the dextran to be homogeneously diluted in the surrounding plasma.

20 As an example of the preferred embodiment, three and eight/tenth (3 and 8/10) milliliters of cryoprotectant is added to the fifteen (15) milliliters of white cells in the white cell bag 30. This addition brings the concentration of DMSO to ten percent (10%) in bag 30.  
25 White cell bag 30 has another outlet 62 which receives a spike 64 from the stem cell freezing bag 60 in an aseptic manner. The white cell bag 30 communicates with stem cell freezing bag 60 via conduit 66 controlled by clamp 67. The cryoprotected white cell mixture is received into the  
30 stem cell freezing bag 60 via portal 68. The portal 68 is specially configured to include a stand tube which allows a standing column of stem cell mixture to be retained therewithin for sequestering into a series of compartments 70, each spaced from the other by heat seals 72. These  
35 specimens 70 can be used for pre-infusion confirmation of an optimum HLA match or other tests, once a particular stem cell freezing bag 60 has been chosen as appropriate

-15-

for the putative recipient.

The stem cell freezing bag 60 is further characterized by having a plurality of compartments within the main body of the bag 60, each compartment provided with indicia 75 thereon for identification of the specific unit, establishing a form of chain of custody. More particularly, the stem cell freezing bag 60 includes at least a first major portion 74 and a second minor portion 76. Typically, the ratio between the major portion 74 and the minor portion 76 is eighty percent (80%) major portion and twenty percent (20%) minor portion. These portions of bag 60 are delimited by heat seal 80 and, after filling, contribute to dividing the stem cell freezing bag into two, intimately attached, but independent white cell containers once heat seals at both locations 82 are executed.

Each portion is in communication with its own outlet. The major portion 74 is in communication with its portal 84 while the minor portion 76 communicates with its own portal 86. In addition, the heat seal location may include a line of demarcation 81 defining a scoreline which allows the major portion 74 to be severed, without thawing, from the minor portion 76. It is contemplated that the stem cells contained in the minor portion 76 can be allocated for other uses, such as for increasing the numbers of useful cells by culturing the stem and progenitor cells in a propagation medium. The stem cells in major portion 74 are left undisturbed for administration as transplants. The freezer bag 60 and stand tube/portal 68 have negligible thickness. The purpose of this particular geometry is to assure that the white cells in compartments 76, 74 and 70 all maintain a uniform and narrow thickness so that subsequent freezing regimens achieve near identical controlled rate freezing conditions.

In a preferred embodiment, approximately nineteen (19) milliliters of therapeutic product are

-16-

contained within the freezing bag 60. The stem cell freezing bag 60 is gradually frozen to an extremely low temperature such as in liquid nitrogen for permanent storage. This preserves the stem cells in a state such that, upon thawing, they are recovered in quantity and exhibit a high degree of cell viability.

Once it has been determined that the given stem cells within a freezing bag 60 are to be used in a transplant procedure, the stem cells are first thawed to a temperature where the stem cells and constituent components change phase back from a solid to a liquid. Next, the stem cells are washed to remove the cryoprotectant which was added prior to freezing. Preferably, the wash is intended to remove the DMSO by using an isotonic fluid, preferably a colloid. For example, a mixture having five percent (5%) albumin and ten percent (10%) dextran in a saline solution is used to dilute the DMSO in the extracellular environment and secondarily reduce its concentration inside the white blood cells. Subsequently, the mixture is centrifuged at four hundred (400) gs for ten (10) minutes with the supernatant expressed therefrom.

As mentioned supra, the enriched white cells were present in volume at approximately fifteen (15) milliliters prior to the addition of three point eight (3.8) milliliters of cryoprotectant. When placed in the stem cell freezing bag, about four (4) milliliters were placed in the secondary compartment 76 and fifteen (15) milliliters were retained in the primary container 74. In actuality, somewhat less than the four (4) milliliters are allocated as is just described because the stem cell samples contained within compartments 70 may contain collectively up to one (1) milliliter. In any event, the thawed white blood and stem cell suspension prior to washing contained ten (10%) cryoprotectant by volume. After the dilution, spinning and expressing off the supernatant the sedimented stem cells (typically in a

-17-

volume less than three (3) milliliters) are diluted once again to a volume adequate for administration to the recipient, fifteen (15) milliliters or more. This second dilution reduces the concentration of DMSO to below one percent (1%). Therefore, the quantity of DMSO retained is in the order to one-tenth (1/10) gram. This is very much less, compared with the prior art which typically may have involved two hundred (200) milliliters of ten percent (10%) DMSO i.e. twenty (20) grams of this compound.

In addition, the therapeutic dose described in the disclosed invention hereinabove has a special efficacy because the processing described hereinabove has removed from the whole blood, the bulk of the red cells, plasma, cryoprotectant, free hemoglobin, etc. which heretofore have exhibited adverse consequences on the recipient and has restored the osmolarity of the stem and progenitor cells to the normal range of three hundred (300) milliosmols from the over one thousand (1000) milliosmols of ten percent (10%) DMSO solution.

It is to be noted that the stem cells that are stored in freezing bags must be kept at extremely low temperatures such as those achievable using liquid nitrogen. By providing white stem cells in twenty (20) milliliter quantities, the problems that would have existed before in the provision of storage space for units with ten fold larger volumes of cryopreserved placental blood (whole) will have been solved by the smaller storage requirement of separated white blood cells associated with the instant invention.

One attribute of the instant invention is that the therapeutic dose involves a relatively low level of DMSO in the finished product that is to be administered. A second attribute involves the fact that a ten (10) fold lower concentration of red blood cells are contained in a unit without significant loss of stem and progenitor cells. The lower red blood cell numbers reduce the presence of hemoglobin in the thawed specimen and decrease

-18-

the problems associated with red blood cell incompatibilities. Further, the viability of the white cells contained in the dose after thawing is typically three (3) to four (4) fold higher than in the prior art, particularly after administration and dilution in the recipient's own plasma. Experimentally, thawed white cells are diluted in twenty (20) milliliters of plasma prior to counting for viability. In prior art, unwashed white cell viability was typically of the order of twenty percent (20%). According to the present invention, using the DNA fluorescence stain or other viability tests, the mononuclear cells are much greater than twenty percent (20%), typically greater than ninety percent (90%) viable. When stem and progenitor cells are cultured in vitro from such white cell concentrates after thawing as described, the number of viable cells estimated by the number of colonies formed is also greater than ninety percent (90%) of the original numbers.

While the previous discussion has focused on the desirability of using cord blood from placental stem cells, other peripheral stem cells can also be processed in a similar manner to provide benefits. Further, having thus described the invention it should be apparent that numerous structural modification and adaptations of the bag set, the chemical nature of the reagents and cryoprotectants and the details of the processing steps, may be resorted to without departing from the scope and fair meaning of the instant invention as set forth hereinabove and as described hereinbelow by the claims.

30



-19-

What is Claimed is:

1. A system for concentrating blood cells, comprising in combination:

5 a first blood bag adapted to receive blood from a source,

means within said blood bag to prevent coagulation,

reagent means removably coupled to said blood bag,

10 means for separating white blood cells containing plasma supernatant from said first blood bag and into a satellite white cell bag,

means for separating white cells from the plasma in said white cell bag, and

15 a satellite plasma bag removably coupled to said white cell bag for receiving the expressed supernatant plasma from said white cell bag.

20 2. The system of claim 1 further including cryoprotectant means operatively coupled to said white cell bag.

3. The system of claim 2 further including a stem cell freezing bag operatively coupled to said white cell bag for transferring contents from said white cell bag to said stem cell freezing bag.

25 4. The system of claim 3 wherein said stem cell freezing bag includes a plurality of compartments therewithin.

30 5. The system of claim 4 wherein said plurality of compartments include means for sealing said compartments into isolated sequestered areas and means on said sequestered areas for indicating the origin of said freezing bag partitioned area.

35 6. The system of claim 5 including means for washing cryoprotectant from said contents of said freezing bag.

-20-

7. The system of claim 6 further including means for metering said cryoprotectant into said white cell bag.

8. The system of claim 7 further including  
5 clamping means to constrict passage from one bag to another.

9. A method for concentrating placental stem cells, comprising the steps of:

10 placing blood into a first blood bag,  
preventing coagulation within said blood bag,  
coupling reagent means into said blood bag,  
separating supernatant from said first blood bag  
and placing the supernatant into a white cell bag,  
separating white cells from plasma in said white  
15 cell bag, and

removably coupling a plasma bag to said white cell bag and expressing the plasma from the white cell bag into said plasma bag.

10. The method of claim 9 further including  
20 coupling cryoprotectant means to said white cell bag.

11. The method of claim 10 further including transferring contents from said white cell bag to a stem cell freezing bag, and freezing the contents of the stem cell freezing bag.

25 12. The method of claim 11 further including sequestering the contents in said stem cell freezing bag into a plurality of partitioned areas prior to freezing the contents.

30 13. The method of claim 12 further including labeling the partitioned areas of the stem cell freezing bag to identify the source of the cells contained therewithin.

35 14. The method of claim 13 further including removing one of the partitions from the stem cell freezing bag to confirm the contents.

-21-

15. The method of claim 14 including thawing the contents of the stem cell freezing bag upon ascertaining the matching of that bag.

5 16. The method of claim 15 including the step of washing cryoprotectant from the contents of one partitioned area of the stem cell freezing bag.

17. The method of claim 16 including the step of diluting the washed contents of the stem cell freezing bag and injecting the contents into a patient.

10 18. A therapeutic product comprising white blood cells in a solution having a constituent presence of at least one white cell part per one hundred red cells and a stem and progenitor cell viability greater than 80%.

15 19. The product of claim 18 further comprising said white cells in a solution having an intracellular cryoprotectant concentration below 1%.

20. The product of claim 18 in which the stem and progenitor cells have an osmolarity in the range of three hundred milliosmols.

20 21. The product of claim 20 wherein the prescribed dose of said product is in the range of ten to twenty milliliters.

22. The product of claim 21 where the stem and progenitor cells are present in the ratio of approximately 25 90% with respect to the original donated blood.

23. The product of 18 wherein there is a reduction of 90% of the red cells from an original blood donation.

24. A therapeutic product comprising white 30 blood cells in a solution having a constituent presence of at least one white blood cell per one hundred red blood cells formed from the process of:

35 placing blood into a first blood bag,  
preventing coagulation within said blood bag,  
coupling reagent means into said blood bag,  
separating supernatant from said first blood bag  
and placing the supernatant into a white cell bag,

-22-

separating white cells from plasma in said white cell bag,

removably coupling a plasma bag to said white cell bag and expressing the plasma from the white cell bag into said plasma bag,

coupling cryoprotectant means to said white cell bag,

transferring contents from said white cell bag to a stem cell freezing bag,

freezing the contents of the stem cell freezing bag,

sequestering the contents in said stem cell freezing bag into a plurality of partitioned areas prior to freezing the contents,

labeling the partitioned areas of the stem cell freezing bag to identify the source of the cells contained therewithin,

removing one of the partitions from the stem cell freezing bag to confirm the contents,

thawing the contents of the stem cell freezing bag upon ascertaining the matching of that bag,

washing cryoprotectant from the contents of one partitioned area of the stem cell freezing bag, and

diluting the washed contents of the stem cell freezing bag and injecting the contents into a patient.

25. A method for removing at least 90% of red blood cells and plasma from placental, umbilical cord and/or peripheral donor blood while retaining a composition comprising hematopoietic stem and progenitor cells having a viability of at least 80%, said method comprising performing the following steps in an integrated, aseptic system:

contacting placental, umbilical cord and/or peripheral donor blood with an anticoagulant to prevent coagulation of said donor blood;

contacting said anticoagulated blood with a rouleaux-inducing reagent to facilitate at least 90% of

-23-

red blood cells contained in said donor blood to form a rouleaux;

gently centrifuging said rouleaux-inducing reagent contacted blood to obtain a first sediment and a first supernatant, wherein said first sediment comprises at least 90% of red blood cells contained in said donor blood;

isolating said first supernatant;

centrifuging said first supernatant to obtain a second sediment and a second supernatant, wherein said second supernatant comprises at least 90% of plasma contained in said donor blood; and

removing said second supernatant so as to retain a composition comprising hematopoietic stem and progenitor cells having a viability of at least 80%, and less than 10% of red blood cells and plasma contained in said donor blood.

26. The method of claim 25 wherein said integrated, aseptic system comprises a series of interconnected bags.

27. The method of claim 25 wherein said anticoagulant comprises citrate, phosphate and dextrose.

28. The method of claim 25 wherein said rouleaux-inducing reagent comprises hydroxyethyl starch.

29. The method of claim 25 which further comprises contacting said composition with a cryoprotectant at a rate and an amount sufficient to obtain a cryoprotected composition without damaging cells contained in said composition; and freezing said cryoprotected composition.

30. The method of claim 29 wherein said cryoprotectant comprises dimethylsulfoxide.

31. The method of claim 30 wherein said cryoprotectant further comprises dextran.

32. The method of claim 30 wherein the final concentration of dimethylsulfoxide in the cryoprotected suspension is 10%.

-24-

33. A method for removing at least 90% of red blood cells and plasma from placental, umbilical cord and/or peripheral donor blood while retaining a composition comprising hematopoietic stem and progenitor cells having a viability of at least 80%, said method comprising performing the following steps in a series of interconnected bags under aseptic conditions:

contacting placental, umbilical cord and/or peripheral donor blood in a blood bag with an anticoagulant to prevent coagulation of said donor blood;

contacting said anticoagulated blood in said blood bag with a rouleaux-inducing reagent to facilitate at least 90% of red blood cells contained in said donor blood to form a rouleaux;

gently centrifuging said blood bag to obtain a first sediment and a first supernatant, wherein said first sediment comprises at least 90% of red blood cells contained in said donor blood;

expressing said first supernatant into a white cell bag;

centrifuging said white cell bag to obtain a second sediment and a second supernatant, wherein said second supernatant comprises at least 90% of plasma contained in said donor blood; and

expressing said second supernatant into a plasma bag so as to retain a composition in said white cell bag comprising hematopoietic stem and progenitor cells having a viability of at least 80%, and less than 10% of red blood cells and plasma contained in said donor blood.

34. The method of claim 33 wherein said anticoagulant comprises citrate, phosphate and dextrose.

35. The method of claim 33 wherein said rouleaux-inducing reagent comprises hydroxyethyl starch.

36. The method of claim 33 which further comprises contacting said composition with a cryoprotectant at a rate and an amount sufficient to obtain a cryoprotected composition without damaging cells

-25-

contained in said composition; and freezing said cryoprotected composition.

37. The method of claim 36 wherein said cryoprotectant comprises dimethylsulfoxide.

5           38. The method of claim 37 wherein said cryoprotectant further comprises dextran.

39. The method of claim 38 wherein the final concentration of dimethylsulfoxide in the cryoprotected suspension is 10%.

10

1/3




Figure 1



2/3




Figure 2





Figure 3

3/3

Figure 4



Europäisches  
Patentamt

# EUROPÄISCHER RECHERCHENBERICHT

Nummer der Anmeldung  
EP 97 12 2072

EINSCHLÄGIGE DOKUMENTE			
Kategorie	Kennzeichnung des Dokuments mit Angabe, soweit erforderlich, der maßgeblichen Teile	Betrifft Anspruch	KLASSIFIKATION DER ANMELDUNG (Int.Cl.6)
X	US 4 731 330 A (HILL JAMES L ET AL) * Ansprüche * * Spalte 2, Zeile 13 - Zeile 17 * * Spalte 3, Zeile 4 - Zeile 9 * * Spalte 6, Zeile 41 - Spalte 7, Zeile 31 * ---	1-9	G01N33/96 G01N33/49
X	WO 96 17514 A (NEW YORK BLOOD CENTER INC) * Ansprüche * * Seite 16, Zeile 23 - Seite 17, Zeile 29 * ---	2-5	
A	US 4 701 417 A (PORTENHAUSER RUDOLF ET AL) * das ganze Dokument * ---	1-9	
A	US 4 219 440 A (ARMSTRONG DOUGLAS ET AL) * das ganze Dokument * ---	1-9	
A	FR 2 475 737 A (PASTEUR INSTITUT) * das ganze Dokument * -----	1-9	
			RECHERCHIERTE SACHGEBIETE (Int.Cl.6)
			G01N
Der vorliegende Recherchenbericht wurde für alle Patentansprüche erstellt			
Recherchenort DEN HAAG		Abschlußdatum der Recherche 24. März 1998	Prüfer Routledge, B
KATEGORIE DER GENANNTEN DOKUMENTE			
X : von besonderer Bedeutung allein betrachtet Y : von besonderer Bedeutung in Verbindung mit einer anderen Veröffentlichung derselben Kategorie A : technologischer Hintergrund O : nichtschriftliche Offenbarung P : Zwischenliteratur		T : der Erfindung zugrunde liegende Theorien oder Grundsätze E : älteres Patentedokument, das jedoch erst am oder nach dem Anmeldedatum veröffentlicht worden ist D : in der Anmeldung angeführtes Dokument L : aus anderen Gründen angeführtes Dokument & : Mitglied der gleichen Patentfamilie, übereinstimmendes Dokument	

**ANHANG ZUM EUROPÄISCHEN RECHERCHENBERICHT  
ÜBER DIE EUROPÄISCHE PATENTANMELDUNG NR.**

EP 97 12 2072

In diesem Anhang sind die Mitglieder der Patentfamilien der im obengenannten europäischen Recherchenbericht angeführten Patentdokumente angegeben.

Die Angaben über die Familienmitglieder entsprechen dem Stand der Datei des Europäischen Patentamts am  
Diese Angaben dienen nur zur Unterrichtung und erfolgen ohne Gewähr.

24-03-1998

Im Recherchenbericht angeführtes Patentdokument	Datum der Veröffentlichung	Mitglied(er) der Patentfamilie	Datum der Veröffentlichung
US 4731330 A	15-03-88	KEINE	
WO 9617514 A	13-06-96	AU 4507896 A CA 2207044 A EP 0796040 A	26-06-96 13-06-96 24-09-97
US 4701417 A	20-10-87	DE 3445010 A EP 0184765 A JP 4071468 B JP 61142467 A	19-06-86 18-06-86 13-11-92 30-06-86
US 4219440 A	26-08-80	AU 525932 B AU 4806779 A CA 1120835 A DE 2923957 A DK 248679 A FR 2432172 A GB 2023287 A,B NL 7904656 A SE 7905238 A	09-12-82 20-12-79 30-03-82 03-01-80 16-12-79 22-02-80 28-12-79 18-12-79 16-12-79
FR 2475737 A	14-08-81	KEINE	